

Date: 4<sup>th</sup> November 2016  
Stud: Noorilim  
Horse: Glacier Bay 16

Your horse has had a transphyseal screw placed in the left fore distal radius and right fore cannon

Your horse requires;

- 8mls penicillin in the muscle twice daily for 3 days.
- Confinement in a yard for two weeks,
- Bandaging of the limb for 2 weeks, changing the bandage every 4 days
- Sutures should be removed 12-14 days after surgery; please take care to cut one side of the loop only, and pull the whole stitch out

Please return to the hospital for screw removal when the limb straightens. Some swelling around the surgical site is expected and usually resolves after screw removal

*Please call immediately if concerned; particularly if the horse is dull or off its feed or lame*

Please call to discuss progress at any point,

Sarah Jalim, BVM&S, FACVSc  
Registered Specialist in Equine Surgery  
0409 977518

Sarah L Jalim  
BVM&S FACVSc  
Specialist in Equine Surgery

Simon J Robinson  
BVSc BScAgr MACVSc Dip ACT  
(Registered Specialist in Equine Reproduction)

Michael J Whiteford  
BVMS MRCV

Tom Russell  
BVMS MACVs Dip ECVS  
(Registered Specialist in Equine Surgery)

**Bendigo Equine Hospital**  
38 Heinz Street, White Hills, Vic, 3550  
T: (03) 5448 5331 F: (03) 5448 5938

**Victorian Equine Reproduction Centre & Vic Equine**  
1000 Toolamba Road, Toolamba Vic 3614  
T: 0400 504 497 F: (03) 5835 6477