

**Horsemed SA - Morphettville Equine
Clinic**

90 Morphett Road,
Glengowrie, SA 5044
Ph: 08 8294 8900
Fax: 08 8376 0950
Email: admin@horsemedsa.com
ABN 17 126 239 530



**Horsemed SA - Hills Equine
Clinic**

43 Old Mount Barker Road
(PO Box 538)
Echunga SA 5153
Ph: 08 8391 9400
Email: hills@horsemedsa.com
www.horsemedsa.com

MAGIC MARINER

Microchip: 985100012210886

Trainer: Andrew Gluyas

Examination date: 16/6/2025

The gelding was noticed to have some superficial swelling to the right fore mid tendon over the course of the previous week. The swelling did subside on its own accord but the horse was found to be slightly lame on the right fore when trotted up.

Palpation of the origin of the suspensory ligament in the right fore did illicit a pain response. The suspensory was scanned and a small anechoic lesion was found at the bifurcation of the suspensory. It is low grade.

Treatment options include 1: rest alone, and possible re-scan in 6 months before returning to work.

Option 2: treat the area with extra corporal shockwave therapy (ECSWT) – this would entail three treatments two weeks apart. The first treatment can be done here in the stables and the follow ups if required could be done where ever the horse is spelled. The advantage of treating with shock wave is that the rehabilitation period is reduced and he can come back into work sooner – usually after approximately three months.

The lesion is small and his future as a racehorse would be considered good to excellent.

Regards,

Dr. Andrew Hunt